arXiv:1406.0411v1 [math.FA] 2 Jun 2014

# A NEW PROOF FOR THE BORNOLOGICITY OF THE SPACE OF SLOWLY INCREASING FUNCTIONS

JULIAN LARCHER AND JOCHEN WENGENROTH

Abstract. A. Grothendieck proved at the end of his thesis that the space $\mathcal{O}_{\mathrm{M}}$ of slowly increasing functions and the space $\mathcal{O}'_C$ of rapidly decreasing distributions are bornological. Grothendieck's proof relies on the isomorphy of these spaces to a sequence space and we present the first proof that does not utilize this fact by using homological methods and, in particular, the derived projective limit functor.

## 1. Introduction and notation

In [Sch66, p. 243] L. Schwartz introduced the space of multipliers of temperate distributions, i.e., the space of slowly increasing functions

$$\mathcal{O}_{\mathrm{M}} = \{f \in \mathcal{C}^\infty(\mathbb{R}^d)\,;\, \forall \alpha \in \mathbb{N}_0^d\ \exists N \in \mathbb{N} :\ \langle x\rangle^{-N}\partial^\alpha f \in L^\infty\},$$

where $\mathcal{C}^\infty(\mathbb{R}^d)$ is the space of complex valued, infinitely differentiable functions on $\mathbb{R}^d$, $\langle x\rangle = 1 + |x|^2$, $\partial^\alpha$ is the partial derivative, and $L^\infty$ is the Lebesgue space of bounded functions. The dual $\mathcal{O}'_M$ of $\mathcal{O}_{\mathrm{M}}$ is the space of very rapidly decreasing distributions.

Schwartz also introduced the space of convolutors of temperate distributions, i.e., the space $\mathcal{O}'_C$ of rapidly decreasing distributions, which is the dual of the space

$$\mathcal{O}_{\mathrm{C}} = \{f \in \mathcal{C}^\infty(\mathbb{R}^d)\,;\, \exists N\, \forall \alpha \in \mathbb{N}_0^d :\ \langle x\rangle^{-N}\partial^\alpha f \in L^\infty\}$$

of very slowly increasing functions. These spaces are related as in the diagram

$$\begin{array}{ccc} \mathcal{O}_{\mathrm{C}} & \subseteq & \mathcal{O}_{\mathrm{M}} \\ \wr\| & & \wr\| \\ \mathcal{O}'_M & \subseteq & \mathcal{O}'_C \end{array}$$

where in both cases the Fourier transform can be taken as the isomorphism.

It is comparatively easy to see that the four spaces are nuclear and semi-reflexive, that $\mathcal{O}_{\mathrm{M}}$ and $\mathcal{O}'_C$ are complete and that $\mathcal{O}_{\mathrm{C}}$ and $\mathcal{O}'_M$ are (LF)-spaces and hence bornological. But the completeness of $\mathcal{O}_{\mathrm{C}}$ and $\mathcal{O}'_M$ and the bornologicity of $\mathcal{O}_{\mathrm{M}}$ and $\mathcal{O}'_C$ are not trivial (which was even asserted by Grothendieck, [Gro55, Chap. II, p. 130]). Since the dual of a bornological space is complete and the dual of a complete nuclear space is bornological, these two problems are equivalent (for the definitions of these topological properties and relations between them see [Itō87, Section 424]).

Grothendieck proved that $\mathcal{O}_{\mathrm{M}}$ is bornological by showing that it is isomorphic to a complemented subspace of the sequence space $s\,\hat{\otimes}_\pi\, s'$ [Gro55, Chap. II, Lemme 18, p. 132] and verified "directly" that the space $s\,\hat{\otimes}_\pi\, s'$ is bornological [Gro55, Chap.II, Prop. 15, p. 125, Cor. 2, p. 128]. We will find out more about this isomorphy in

2000 *Mathematics Subject Classification.* Primary 46M18 · 46F05 ; Secondary 46A08 · 46A45.
*Key words and phrases.* derived projective limit functor · slowly increasing functions · rapidly decreasing distributions · sequence space representations.

Section 2 and also give a homological proof of the bornologicity of $s\,\hat{\otimes}_\pi\, s'$.

In [Kuc85], J. Kučera claimed to have presented a new (and simple) proof for the main properties of the space $\mathcal{O}_{\mathrm{M}}$. That Kučera's proof contains severe mistakes and that it is based on incorrect propositions is clarified in [Lar12], where also the lack of a proof of the bornologicity of $\mathcal{O}_{\mathrm{M}}$, that does not use the isomorphy $\mathcal{O}_{\mathrm{M}} \cong s\,\hat{\otimes}_\pi\, s'$, is pointed out. In Section 3 we will give such a proof.

## 2. Projective limits and the space $s\,\hat{\otimes}_\pi\, s'$

Since quotients (and, in particular, complemented subspaces) of bornological spaces are bornological, it was sufficient for Grothendieck to prove that $\mathcal{O}_{\mathrm{M}}$ is isomorphic to a complemented subspace of $s\,\hat{\otimes}_\pi\, s'$, where $s$ is the space of rapidly decreasing sequences

$$s = \{(x_j)_{j\in\mathbb{N}} \in \mathbb{C}^{\mathbb{N}}\,;\, \forall k : \sup_{j\in\mathbb{N}} j^k|x_j| < \infty\}$$

and $s'$ is its dual, the space of slowly increasing sequences

$$s' = \{(x_j)_{j\in\mathbb{N}} \in \mathbb{C}^{\mathbb{N}}\,;\, \exists k : \sup_{j\in\mathbb{N}} j^{-k}|x_j| < \infty\}.$$

By $s\otimes_\pi s'$ we denote the completed projective tensor product of these spaces. E.g., by [Bar12, Remark 1, p. 321], this space $s\,\hat{\otimes}_\pi\, s'$ is canonically isomorphic to

$$s\,\hat{\otimes}_\pi\, s' \cong \{x \in \mathbb{C}^{\mathbb{N}\times\mathbb{N}}\,;\, \forall n\,\exists N : \sup_{i,j} i^n j^{-N}|x_{i,j}| < \infty\}.$$

In [Val81], M. Valdivia proved that $\mathcal{O}_{\mathrm{M}}$ is even isomorphic to $s\,\hat{\otimes}_\pi\, s'$ itself which answered a question posed in [Gro55, Chap. II, p. 134]. C. Bargetz used this fact, the bornologicity of $s\,\hat{\otimes}_\pi\, s'$, and methods of the theory of topological tensor products to obtain the isomorphy $\mathcal{O}_{\mathrm{C}} \cong s\hat{\otimes}_\iota s'$ [Bar12, Prop. 1, p. 318].

The descriptions of the spaces $\mathcal{O}_{\mathrm{M}}$ and $s\hat{\otimes}_\pi s'$ already indicate how they can be written as projective limits of LB-spaces (countable inductive limits of Banach spaces)

$$\mathcal{O}_{\mathrm{M}} = \bigcap_{n\in\mathbb{N}} X_n = \bigcap_{n\in\mathbb{N}} \bigcup_{N\in\mathbb{N}} X_{n,N}, \tag{1}$$

$$s\hat{\otimes}_\pi s' = \bigcap_{n\in\mathbb{N}} Y_n = \bigcap_{n\in\mathbb{N}} \bigcup_{N\in\mathbb{N}} Y_{n,N}, \tag{2}$$

where $X_{n,N}$ and $Y_{n,N}$ are the Banach spaces

$$X_{n,N} = \{f \in \mathcal{C}^n(\mathbb{R}^d)\,;\, \|f\|_{n,N} = \sup_{x\in\mathbb{R}^d,|\alpha|\le n} \langle x\rangle^{-N}|\partial^\alpha f(x)| < \infty\},$$

$$Y_{n,N} = \{x \in \mathbb{C}^{\mathbb{N}\times\mathbb{N}}\,;\, \|x\|_{n,N} = \sup_{i,j} i^n j^{-N}|x_{i,j}| < \infty\}.$$

These representations as projective limits of LB-spaces are not only natural but also extremely useful since there are very good criteria for checking bornologicity. They are related to the derived projective limit functor $\mathrm{Proj}^1\mathscr{X}$ (which can be defined as the cokernel of the map $\prod X_n \to \prod X_n$, $(x_n)_n \mapsto (x_n - \varrho^n_{n+1}(x_{n+1}))_n$ where $\varrho^n_m$ are the connecting maps of the projective spectrum $\mathscr{X}$, in our cases, $\varrho^n_m$ are just inclusions). Indeed, an unbublished theorem of D. Vogt (his proof reproduced in [Wen03, Th. 3.3.4]) says that $\mathrm{Proj}\,\mathscr{X}$ is bornological whenever $\mathrm{Proj}^1\,\mathscr{X} = 0$. Moreover, there is a variety of evaluable conditions ensuring $\mathrm{Proj}^1\,\mathscr{X} = 0$. We are going to apply the following results of Palamodov-Retakh [Pal71] and the second named author, respectively:

A specturm $\mathscr{X}$ of LB-spaces satisfies $\operatorname{Proj}^1 \mathscr{X} = 0$ if and only if there are Banach discs $D_n$ in $X_n$ with $\varrho^n_m(D_m) \subseteq D_n$ and

$$\forall\, n \in \mathbb{N}\ \exists\, m \geq n\, \forall\, k \geq m:\ \varrho^n_m(X_m) \subseteq \varrho^n_k(X_k) + D_n.$$

The requirement $\varrho^n_m(D_m) \subseteq D_n$ is sometimes very easy to fulfil but in many cases it is very inconvenient. It can be omitted if either all steps $X_n$ are LS-spaces (i.e., the inclusions $X_{n,N} \hookrightarrow X_{n,N+1}$ are compact) or if a slightly stronger condition of Palamodov-Retakh type is required. Denoting by $\varrho^n_\infty : \operatorname{Proj} \mathscr{X} \to X_n$ the obvious map we have:

A specturm $\mathscr{X}$ of LB-spaces satisfies $\operatorname{Proj}^1 \mathscr{X} = 0$ if and only if, for every $n \in \mathbb{N}$, there are a Banach discs $D_n$ in $X_n$ and $m \geq n$ with

$$\varrho^n_m(X_m) \subseteq \varrho^n_\infty(\operatorname{Proj} \mathscr{X}) + D_n.$$

We refer to [Wen03] for the proofs of these characterization and much more information about derived functors. Typically, the decompositions required in conditions of Retakh-Palamodov type are quite easy to produce in the case of spaces of sequences (or matrices) since one can write $x = \chi\, x + (1-\chi)x$ where $\chi$ is the indicator function of a suitably chosen set. We want to exemplify this by giving a very short proof for the bornologicity of $s\hat{\otimes}_\pi s'$ (which is similar to Vogt's proof of $\operatorname{Ext}^1(s,s) = 0$ [Vog84, Lemma 2.1, p. 359]).

**Proposition 1.** *The space $s\hat{\otimes}_\pi s'$ is bornological.*

*Proof.* We keep the notation $s\hat{\otimes}_\pi s' \cong \bigcap_{n\in\mathbb{N}} Y_n = \bigcap_{n\in\mathbb{N}} \bigcup_{N\in\mathbb{N}} Y_{n,N}$ from above and we will verify the Palamodov-Retakh condition for the unit balls $D_n$ of $Y_{n,0}$ which trivially satisfy $D_{n+1} \subseteq D_n$. For $n \in \mathbb{N}$ we take $m = n+1$ and fix $x \in Y_n$ as well as $k \geq n+1$. Since $x \in Y_{m,M}$ for some $M \in \mathbb{N}$ we have

$$\|x\|_{m,M} = \sup_{i,j} i^m j^{-N} |x_{i,j}| = c < \infty.$$

We set $y_{i,j} = x_{i,j}$ if $i < cj^M$ and $y_{i,j} = 0$ else, as well as $z = x - y$. For $i < cj^m$ we have $z_{i,j} = 0$ and for $i \geq cj^M$ we estimate

$$i^n j^{-0} |z_{i,j}| = i^m j^{-M} |z_{i,j}|\, j^M / i \leq \|x\|_{m,M}/c = 1$$

which proves $z \in D_n$. It remains to show $y \in Y_{k,K}$ for $K$ sufficiently large. Indeed, for $K = M(k-m+1)$ we have $y_{i,j} = 0$ if $i \geq cj^M$ and if $i < cj^M$ we estimate

$$i^k j^{-K} |y_{i,j}| = i^m j^{-M} |y_{i,j}| i^{k-m} j^{M-K} \leq \|x\|_{m,M} c^{k-m} j^{(k-m)M+M-K} = c^{k-m+1}.$$

This proves $\|y\|_{k,K} < \infty$, as required. □

## 3. The new proof

Now we want to prove $\operatorname{Proj}^1 \mathscr{X} = 0$ for the spectrum $\mathscr{X} = (X_n)_{n\in\mathbb{N}}$ in (1) in order to obtain that $\mathcal{O}_{\mathrm{M}}$ is bornological. Splitting up a given function $f \in X_m$ as $f = \chi f + (1-\chi) f$ with a cut-off function $\chi$ (as in the proof of Proposition 1) does not work in this case. But we will see how $f$ can be "split up" in the following proof of Grothendieck's result.

**Proposition 2.** *The space $\mathcal{O}_{\mathrm{M}}$ is bornological.*

*Proof.* To obtain $\operatorname{Proj}^1 \mathscr{X} = 0$ we will show

$$\forall n\, \exists m, N:\ X_m \subseteq \mathcal{O}_{\mathrm{M}} + B_{n,N} \tag{3}$$

where $B_{n,N}$ is the unit ball of $X_{n,N}$. This condition means that we have to approximate every $f \in X_m$ with respect to the norm $\|\cdot\|_{n,N}$ by elements of $\mathcal{O}_{\mathrm{M}}$. To achieve such an approximation we use a kernel $K \in \mathcal{O}_{\mathrm{M}}(\mathbb{R}^d \times \mathbb{R}^d)$ satisfying

$$K \geq 0,\ \int_{\mathbb{R}^d} K(t,x)\,dt = 1 \text{ for all } x \in \mathbb{R}^d, \text{ and}$$
$$\operatorname{supp} K(\cdot,x) \subseteq \prod_{j=1}^{d} [x_j, x_j + \varepsilon\langle x\rangle^{-\mu}] =: A_x \text{ for all } x \in \mathbb{R}^d$$

where we will see later how $\varepsilon$ and $\mu$ have to be chosen in dependence on $f \in X_m$. We can obtain such a kernel by defining

$$K(t,x) = \varepsilon^{-d}\langle x\rangle^{\mu d}\varphi(\varepsilon^{-1}\langle x\rangle^{\mu}(t-x))$$

for a positive test function $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ with support in $[0,1]^d$ and $\int_{\mathbb{R}^d}\varphi(t)dt = 1$ (the conditions above can be checked easily and $K \in \mathcal{O}_{\mathrm{M}}$ since every derivative of $K$ can be estimated by a polynomial).

We start with the one-dimensional case $d = 1$ where we can take $m = n+1$ and $N = 0$. So let $f \in X_{n+1,M}$ for some $M \in \mathbb{N}$. We want to find $g \in \mathcal{O}_{\mathrm{M}}$ such that $f - g \in B_{n,0}$. At first we set

$$g_n(x) = \int_{\mathbb{R}} f^{(n)}(t)K(t,x)\,dt$$

and show that this is a good approximation to $f^{(n)}$. Since for $l \in \mathbb{N}_0$

$$\left|g_n^{(l)}(x)\right| = \left|\int_{A_x} f^{(n)}(t)\partial_x^l K(t,x)\,dt\right| \leq \int_{A_x} |P(t)|\,|Q(t,x)|\,dt \leq |R(x)|$$

for some polynomials $P$, $Q$, $R$, the function $g_n$ is contained in $\mathcal{O}_{\mathrm{M}}$. Furthermore we can estimate in virtue of Taylor's formula

$$\left|f^{(n)}(t) - f^{(n)}(x)\right| \leq |t-x|\langle\xi(t,x)\rangle^M \|f\|_{n+1,M}$$

with a point $\xi(t,x)$ between $t$ and $x$. For $\varepsilon$ small enough the inequality $\langle\xi(t,x)\rangle \leq 2\langle x\rangle$ holds for every $x \in \mathbb{R}$ and $t \in A_x$. We obtain

$$\begin{aligned}
|g_n(x) - f^{(n)}(x)| &= \left|\int_{\mathbb{R}} \left(f^{(n)}(t) - f^{(n)}(x)\right) K(t,x)\,dt\right| \\
&\leq \int_{A_x} \left|f^{(n)}(t) - f^{(n)}(x)\right| K(t,x)\,dt \\
&\leq \int_{A_x} |t-x|\langle\xi(t,x)\rangle^M \|f\|_{n+1,M} K(t,x)\,dt \\
&\leq \varepsilon\, 2^M\, \langle x\rangle^{M-\mu}\|f\|_{n+1,M} \int_{A_x} K(t,x)\,dt \\
&= \varepsilon\, 2^M\, \langle x\rangle^{M-\mu}\|f\|_{n+1,M}.
\end{aligned} \tag{4}$$

Now if

$$T : \mathcal{O}_{\mathrm{M}}(\mathbb{R}) \to \mathcal{O}_{\mathrm{M}}(\mathbb{R}), h \mapsto \left(x \mapsto \int_0^x h(t)\,dt\right),$$

we can set

$$g(x) = \sum_{j=0}^{n-1} \frac{f^{(j)}(0)}{j!}x^j + (T^n g_n)(x).$$

Then $g \in \mathcal{O}_{\mathrm{M}}$ and since

$$(T^n f^{(n)})(x) = f(x) - \sum_{j=0}^{n-1} \frac{f^{(j)}(0)}{j!} x^j,$$

integrating (4) (the integral starting at 0) yields

$$|g^{(l)}(x) - f^{(l)}(x)| \leq 1,\ x \in \mathbb{R}^d, l \leq n$$

for $\varepsilon$ small enough and $\mu$ large enough. Hence $g - f \in B_{n,0}$ and the proof is complete for the one-dimensional case.

Now we will prove the two-dimensional case $d = 2$. We set $m = 2n + 1$ and $N = n - 1$ in (3). So let $f \in X_{2n+1,M}$ for some $M$. With the help of a kernel $K \in \mathcal{O}_{\mathrm{M}}(\mathbb{R}^2 \times \mathbb{R}^2)$ like above, we set

$$g_n(x) = \int_{\mathbb{R}^2} \partial^{(n,n)} f(t) K(t,x)\, dt$$

in order to approximate $\partial^{(n,n)} f$ by $g_n$. Similar to the one-dimensional case we have

$$\left|\partial^{(n,n)} f(t) - \partial^{(n,n)} f(x)\right| \leq c \cdot |t - x| \langle \xi(t,x) \rangle^M \|f\|_{2n+1,M}$$

and $\langle \xi(t,x) \rangle \leq 2\langle x \rangle$ for $t \in A_x$ and $\varepsilon$ small enough and thus

$$\begin{aligned} |g_n(x) - \partial^{(n,n)} f(x)| &\leq c \int_{A_x} |t - x| \langle \xi(t,x) \rangle^M \|f\|_{2n+1,M} K(t,x)\, dt \\ &\leq \tilde{c}\, \varepsilon\, \langle x \rangle^{M-\mu} \|f\|_{2n+1,M}. \end{aligned} \tag{5}$$

Let us denote $T_j$ the integral with respect to the $j$-th component (the integral starting at 0). Applying $T_1 \circ T_2$ $n$-times to $\partial^{(n,n)} f(x)$ yields

$$(T_1^n T_2^n f)(x) = f(x) + \sum_{\alpha < (n,n)} \partial^\alpha f(0,0) \frac{x^\alpha}{\alpha!} - \sum_{j=0}^{n-1} \partial^{(j,0)} f(0,x_2) \frac{x_1^j}{j!} - \sum_{j=0}^{n-1} \partial^{(0,j)} f(x_1,0) \frac{x_2^j}{j!}.$$

As in the one-dimensional case we can choose $g_0^1, \dots g_{n-1}^1, g_0^2, \dots, g_{n-1}^2 \in \mathcal{O}_{\mathrm{M}}(\mathbb{R})$ such that $\|g_j^1 - \partial^{(0,j)} f(\cdot, 0)\|_{n,0} \leq \varepsilon$ and $\|g_j^2 - \partial^{(j,0)} f(\cdot, 0)\|_{n,0} \leq \varepsilon$. Defining

$$g(x) = (T_1^n T_2^n) g_n(x) - \sum_{\alpha < (n,n)} \partial^\alpha f(0,0) \frac{x^\alpha}{\alpha!} + \sum_{j=0}^{n-1} g_j^2(x_2) \frac{x_1^j}{j!} + \sum_{j=0}^{n-1} g_j^1(x_1) \frac{x_2^j}{j!}$$

and applying $T_1^n T_2^n$ to (5) yields

$$|g(x) - f(x)| \leq \varepsilon + \sum_{j=0}^{n-1} \left( \left|g_j^1(x_1) - \partial^{(0,j)} f(x_1,0)\right| \frac{|x_2|^j}{j!} + \left|g_j^2(x_2) - \partial^{(j,0)} f(0,x_2)\right| \frac{|x_1|^j}{j!} \right)$$

for $\mu$ large enough which implies

$$|g(x) - f(x)| \leq \varepsilon + \varepsilon \sum_{j=0}^{n-1} \frac{|x_2|^j}{j!} + \varepsilon \sum_{j=0}^{n-1} \frac{|x_1|^j}{j!} \leq \varepsilon\, c\, \langle x \rangle^{n-1}$$

for some $c > 1$. Since similar estimates also hold for $|\partial^\alpha g(x) - \partial^\alpha f(x)|$, $|\alpha| \leq n$, we obtain $g - f \in B_{n,n-1}$ and the proof is complete for $d = 2$.

The general case $d \in \mathbb{N}$ is very similar. Inductively we want to show

$$X_{dn+1} \subseteq \mathcal{O}_{\mathrm{M}} + B_{n,(d-1)(n-1)}$$

and start by approximating $\partial^{(n,\dots,n)} f$ by $g_n(x) := \int_{R^d} \partial^{(n,\dots,n)} f(t) K(t,x)\, dt$. Then we integrate the estimate of $g_n - \partial^{(n,\dots,n)} f$ $n$-times with respect to each component. The integral $T_1^n \cdots T_d^n \partial^{(n,\dots,n)} f$ contains $f$ as a summand and terms that are the product of a derivative of $f$ that only depends on less than $d$ components and a polynomial in less than $d$ components with exponents less than $n$. But we can estimate the functions that only depend on less than $d$ variables by the induction hypothesis and hence we can obtain $g \in \mathcal{O}_{\mathrm{M}}$ with $g - f \in B_{n,(d-1)(n-1)}$. □

## References


[]Bar12 Christian Bargetz, *A sequence space representation of L. Schwartz' space $\mathcal{O}_C$*, Arch. Math. (Basel) **98** (2012), no. 4, 317–326. MR 2914348

[]Gro55 Alexander Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **16** (1955). MR 0075539 (17,763c)

[]Itō87 Kiyosi Itō (ed.), *Encyclopedic dictionary of mathematics. Vol. I–IV*, second ed., MIT Press, Cambridge, MA, 1987, Translated from the Japanese. MR 901762 (89b:00033)

[]Kuc85 Jan Kucera, *The spaces $\mathcal{O}_M$ and $\mathcal{O}_C$ are ultrabornological. A new proof*, Internat. J. Math. Math. Sci. **8** (1985), no. 4, 813–816. MR 821639 (87e:46054)

[]Lar12 Julian Larcher, *Some remarks concerning the spaces of multipliers and convolutors, $\mathcal{O}_M$ and $\mathcal{O}'_C$, of Laurent Schwartz*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **106** (2012), no. 2, 407–417. MR 2978922

[]Pal71 Victor P. Palamodov, *Homological methods in the theory of locally convex spaces*, Uspehi Mat. Nauk **26** (1971), no. 1(157), 3–65. MR 0293365 (45 #2442)

[]Sch66 Laurent Schwartz, *Théorie des distributions*, Publications de l'Institut de Mathématique de l'Université de Strasbourg, No. IX–X. Nouvelle édition, entièrement corrigée, refondue et augmentée, Hermann, Paris, 1966. MR 0209834 (35 #730)

[]Val81 Manuel Valdivia, *A representation of the space $\mathcal{O}_M$*, Math. Z. **177** (1981), no. 4, 463–478. MR 624226 (82i:46057)

[]Vog84 Dietmar Vogt, *Some results on continuous linear maps between Fréchet spaces*, Functional analysis: surveys and recent results, III (Paderborn, 1983), North-Holland Math. Stud., vol. 90, North-Holland, Amsterdam, 1984, pp. 349–381. MR 761391 (86i:46075)

[]Wen03 Jochen Wengenroth, *Derived functors in functional analysis*, Lecture Notes in Mathematics, vol. 1810, Springer-Verlag, Berlin, 2003. MR 1977923 (2004d:46090)



Institut für Mathematik, Universität Innsbruck, A-6020 Innsbruck, Austria
*E-mail address*: julian.larcher@uibk.ac.at

Universität Trier, FB IV – Mathematik, 54286 Trier, Germany
*E-mail address*: wengenroth@uni-trier.de